

KARYOTYPE OF THE UKRAINIAN MOUNTAIN-CARPATHIAN SHEEP BREED

V.V. Dzitsiuk^{1*}, K.T. Tipilo¹

¹*Institute of Animal Breeding and Genetics n.d.a. M.V. Zubets of NAAS, Ukraine*

Abstract

The main purpose of this work was to describe the karyotype variability of the Ukrainian Mountain-Carpathian sheep breed. Cytogenetic studies were carried out on a group of 25 sheep Ukrainian Mountain-Carpathian breed. The cultivation of lymphocytes, preparing the cytogenetic samples, classification and registration of chromosome aberrations were held using conventional methods. It was established that the number and structure of the chromosomes of sheep of the Ukrainian Mountain-Carpathian breed corresponds to the norm typical for this species of animals. Chromosomal variability of the studied population includes 19.22% of numerical and structural aberrations. By the number and structure of the chromosome set of the Ukrainian Mountain-Carpathian sheep breed do not differ from other breeds of sheep. Accurate identification of individual chromosomes with routine coloring allows using cytogenetic studies in breeding practice in sheep breeding.

Key words: Ukrainian Mountain-Carpathian breed of sheep, karyotype, chromosomal instability

INTRODUCTION

Animals of the Ukrainian Mountain-Carpathian sheep breed - wool-milk-meat production direction. Their wool is a valuable raw material for both light industry and folk crafts. The local population in the Carpathians is traditionally engaged in the production of Hutsul carpets and bedsteads. From Ukrainian Mountain-Carpathian sheep after weaning lambs for two months of lactation receive 30-40 kg of commodity milk, in the best public utilities and the individual sector - 50 and more kilograms. Milk is used for the production of sheep cheese-brynza, which is in high demand among costumers. Indicators of meat productivity depend on the level of feeding: after feeding on natural mountain pastures, 8-9-month lambs have a live weight of 28-30 kg, after intensive feeding of fattening - 36-38 kg. Slaughter output - 42-45%, the output of flesh from the carcass - 70-75%. After slaughter, the Ukrainian Mountain-Carpathian sheep get excellent quality sheep

skin, which are used for the manufacturing of fur-hood products.

The economic efficiency and profitability of the sheep breeding industry depends on the quality of the product, which is determined by its genetic potential. Therefore, in order to preserve and improve the productive and breeding qualities of animals, one should know the genetic structure of both a separate herd and a breed in general. However, such an important agricultural object as a sheep remains cytogenetically underestimated. The study of the karyotype of sheep has theoretical and practical significance, since in the chromosomes there are genes that determine the development of all the features of the organism and chromosomal aberration of any type can be the reason for reducing their economic value and reproductive function [1-4].

MATERIALS AND METHODS

The object of the study was the number of sheep of the Ukrainian Mountain-Carpathian breed (n=25), which are breeding in the FH "Radwan-Nova" Miloshovichi, Pustomyty district in Lviv region. The cytogenetic study was carried out at the Genetics Laboratory of the Institute of Animal Breeding and Genetics named after MV Zubets (Chubinskoye) using

*Corresponding author:

butterfly221192@gmail.com

The manuscript was received: 26.08.2019

Accepted for publication: 18.12.2019

special techniques and appropriate equipment. For the preparation of chromosomes, samples of culture of lymphocytes of peripheral animal blood were used. Lymphocytes (0,5 ml) were cultured for 72 hours in a nutrient environment RPMI-1640 (2 ml) with the addition of inactivated serum of cattle (0,5 ml), concanavalin (0,1 ml) and gentamicin (0,001 ml – for 1 ml nutrient environment). 2 hours before completion of cultivation, a solution of colchicine was added to the medium at a final concentration of 0,05 $\mu\text{g} / \text{ml}$. The hypotonic treatment was performed using a 0,56 M solution of KCL for 30 minutes, followed by fixation in a freshly prepared and cooled fixative - methyl alcohol and glacial acetic acid (3:1). The routine staining of chromosome preparations was performed by Gimza dye. Cell analysis under a microscope was conducted at an imestion increase of 1000

times and photographed. For analysis and photographing, metaphase plates were selected, in which the chromosomes were separated from each other [5]. The obtained experimental data were worked out by the methods of variation statistics.

RESULTS AND DISCUSSIONS

The cariological analysis carried out by us showed that the diploid set of chromosomes of the sheep of the Ukrainian Mountain-Carpathian breed, as shown by other authors who obtained their results at different times and with the help of cytogenetic technology of various levels [5-7], consists of 54 chromosomes 26 pairs of autosomes and one pair of sex chromosomes (XX or XY) (Fig. 1).




Fig. 1 Preparations of chromosomes at the stage of metaphase: a) chromosomal set in normal; b) aneuploid chromosome set; c) a set with a chromosome with a rupture

The karyotype of the sheep of the Ukrainian Mountain-Carpathian breed consists of chromosomes, which, by morphological structure, can be divided into

two groups: acrocentric and metacentric. In all studied samples, a series of autosomes is represented by three pairs of large metacentrics. The remaining 23 pairs form

a series of gradually decreasing acrocentric chromosomes, most of which have a terminal located centromere. The acrocentric chromosomes of the sheep do not have a significant difference in size, which makes it difficult to identify them without a differentiated color.

The group of metacentric chromosomes distinguishes the first pair, which is obviously larger than other chromosomal pairs. The difference in size between the second and third pairs is less obvious. The ratio of the short shoulder to the long in the first pair is 1:1,3. In the second and third pairs, this ratio is somewhat less. The most successful preparations of the chromosomes of the second and third pairs can be distinguished by the position of centromere, which is located more centrally in chromosomes of the third pair. The remaining autosomes form a series of gradual transitions from large acrocentrics to very small ones.

The individual characteristic of the chromosomes of sheep under routine coloring is very complicated. Most clearly, one can identify the first pair of chromosomes and X-chromosome - the first pair is the largest metacentric, sex X-chromosome - the largest acrocentric. The relative sizes of the X-chromosome considerably vary. Sometimes the predominance of the length of the X-chromosome, comparing to the largest of the acrocentric autosomes, reached 15%, in other chromosomal plates this prevalence was negligible. On separate preparations of the X-chromosome can be isolated according to the elongated short shoulder, the frequency of such morphological features no more than 0,5% (2-3 to 50 examined cells).

It is known from the literature that the Y-chromosome is a small odd sub-metacentric chromosome which present in male karyotypes and absent in females [8]. We examined the karyotype only as a spin, and therefore, the Y chromosome, of course, was not studied.

Chromosomal polymorphism in the form of numerical variations in karyotype (aneuploidy and polyploidy), morphological aberrations, and associations of individual

chromosomes is typical for sheep of the Ukrainian Mountain-Carpathian breed, as well as for other sheep breeds [9, 10]. The results of the analysis of chromosomal variability found that among the 722 analyzed metaphase plates of sheep of the Ukrainian Mountain-Carpathian breed, the proportion of aberrant cells was 19,22%, of which aneuploidy cells – 8,6%, polyploidy – 0,86%, cells with chromosomal ruptures - 2,22%, the frequency of pair chromosomal fragments was 2,66% and the frequency of cells with asynchronous divergence of centromeric regions of chromosomes (ADCRC) was 4,88%.

The main proportion of aneuploids is represented by hypoploids, the proportion of hyperploids, as a rule, is negligible. The stability of the parameters of aneuploidy of the sheep is confirmed by studies of many cytogenetics and can be considered as a special feature. Aneuploidy occurs more often among the sheep than, for example, in cattle and it is formed by the small acrocentrics.

The frequency of polyploidy cells of sheep ranges from 0,53 to 1,36% and is lower than pigs and higher than cattle. The sheep are registered four-, six-, eight-, 16-plodity and more. The bulk of polyploids - tetraploids (64%) and octaploids (21%). The number of triploids and other polyploids is insignificant and is approximately 15%.

In the investigated sheep population, was detected that approximately 2% of the cells have chromosomal ruptures. Structural changes of the chromosomes, in particular translocations, were not detected among animals of the studied population of the Ukrainian Mountain-Carpathian breed.

CONCLUSIONS

By the number and structure of the chromosome set of sheep of the Ukrainian Mountain-Carpathian breed do not differ from other breeds of sheep. From the 27 pairs of chromosomes of the sheep karyotype, the chromosomes of the first group, metacentric structure and chromosomes larger than other, according to size and sex are identified with sufficient accuracy. In connection with this

feature, it is possible to determine the frequency with which these chromosomes participate in structural aberrations, which allows the use of cytogenetic studies in the practice of breeding work in sheep breeding.

REFERENCES

- [1]. Danielak-Czech B., Kaczor U., Sharan M. (2010): Cytomolecular analysis of chromosome X fragility in subfertile sheep. *Animal Biology*, 12(1): 234-239.
- [2]. Brace M.D., Peters O., Menzies P., King W.A., Nino-Soto M.I. (2008): Sex chromosome chimerism and the freemartin syndrome in Rideau Arcott sheep. *Cytogenet Genome Res*, 120: 132–139 <https://doi.org/10.1159/000118752>
- [3]. Iannuzzi A., Perucatti A., Genuardo V., Pauciuolo A., Incarnato D., Musilova P., Rubes J. Iannuzzi C. (2014): The Utility of Chromosome Microdissection in Clinical Cytogenetics: A New Reciprocal Translocation in Sheep. *Cytogenet Genome Res.*, 142(3):174-178. DOI: 10.1159/000361000.
- [4]. Gustavsson I. (1980): Banding techniques in chromosome analysis of domestic animals. *Adv Vet Sci Comp Med.*, 24:245–89.
- [5]. Cribiu E.P., Di Berardino D., Di Meo G.P., Eggen A., Gallagher D.S., Gustavsson I., Hayes H., Iannuzzi L., Popescu C.P., Rubes J., Schmutz S., Stranzinger G., Vaiman A., Womack J. (2001): International System for Chromosome Nomenclature of Domestic Bovids (ISCNDB). *Cytogenet Cell Genet.*, 92: 283-299.
- [6]. Meng L., Gongtao D., Kambol R. (2016): Chromosome karyotype analysis of cultivate somatic cells of Ganjia sheep. *Science Letters.*, 10(1): 8-10.
- [7]. Arslan A., Zima J. (2011): Banded karyotype of the Konya wild sheep (*Ovis orientalis anatolica* Valenciennes, 1856) from Turkey. *Comp. Cytogenet.*, 5: 81-89. Doi: 10.3897/compcytogen.v5i2.1151.
- [8]. Di Meo G.P., Perucatti A., Floriot S. et al. (2005): Chromosome evolution and improved cytogenetic maps of the Y-chromosome in cattle, zebu, river buffalo, sheep and goat. *Chromosome Research*, 13(4): 349-355.
- [9]. Słota E., Bugno-Poniewierska M., Rejduch B., Kozubska-Sobocińska A., Danielak-Czech B. (2007): Molekularne metody w analizach cytogenetycznych genomu zwierząt gospodarskich. *Przegl. Hod.*, 6: 22-25.
- [10]. Dobigny G., Britton-Davidian J., Robinson T.J. (2017): Chromosomal polymorphism in mammals: An evolutionary perspective. *Biological Reviews*, 92(1): 1-22. Doi: 10.1111/brv.12213.